UCSD-PTH-12-10

arXiv:1208.3674v1 [hep-th] 17 Aug 2012

A generalized c-theorem and the consistency of scale without conformal invariance

Jean-François Fortin, Benjamín Grinstein and Andreas Stergiou

Department of Physics, University of California, San Diego, La Jolla, CA 92093 USA

We generalize the proof of the weak version of the c-theorem presented by Komargodski and Schwimmer to the case where a renormalization group flow links a fixed point to a limit cycle, or a cycle to a cycle. We show that the argument of Luty, Polchinski and Rattazzi, ruling out limit cycles in four-dimensional theories is flawed. We compute the S function of Jack and Osborn to third order in the loop expansion and show that it vanishes at fixed points and agrees with the generator Q of limit cycles on them. This vastly simplifies the search for limit cycles, as only one-loop beta functions (two-loop for Yang–Mills couplings) are required to establish their existence.

August 2012

`jffortin@physics.ucsd.edu, bgrinstein@ucsd.edu, stergiou@physics.ucsd.edu`

1. Overview in lieu of Introduction

Two recent, reported results can potentially greatly enrich our understanding of quantum field theory (QFT). On the one hand, Komargodski and Schwimmer (KS) [1], following earlier work by Cappelli, D'Appollonio, Guida and Magnoli (CDGM) [2], have delineated a proof, without resource to perturbation theory, of an inequality satisfied when a four-dimensional QFT flows between two fixed points of the renormalization group (RG). On the other hand, we have discovered closed renormalization group trajectories in four-dimensional theories, in a regime where perturbation theory is applicable [3–6]. While the former result can impose restrictions on the possible realizations of long distance (IR) phases of QFTs, the latter gives explicitly new unexplored phases of QFTs. A question naturally arises as to whether these results are compatible.

In this work we extend the work of CDGM and KS to include cycles.[1] More concretely, KS assume the existence of a flow from a short distance (UV) fixed point to an IR fixed point, and argue that the coefficient of the Euler density in the trace anomaly, denoted by a, is larger at the UV than the IR fixed points: $a_{\mathrm{UV}} > a_{\mathrm{IR}}$. This, then, is a proof of the "weak version of the c-theorem." We will show that we can extend the argument to include putative flows from a fixed point or cycle to another fixed point or cycle. Along their argument KS make several explicit and implicit assumptions. We make the same assumptions, and we try to be as explicit as possible about them. We make no additional assumptions.

Recently, Luty, Polchinski and Rattazzi (LPR) have extended the KS argument to argue, to lowest order in perturbation theory, that a decreases monotonically along the RG-flow towards the IR [7]. This would then be a proof of the "strong version of the c-theorem." Additionally, LPR compute the rate of change of a as it flows along a cycle and show that a decreases monotonically in that case as well. Were their results correct they would render our results invalid: LPR argue that the monotonic decrease of a along flows on cycles rules out their existence in four-dimensional relativistic field theories.

However, the computations presented by LPR are interpreted by them incorrectly. This is easy to see by the known fact that a increases away from trivial UV fixed points: for a Yang–Mills theory (YM) with beta function $\beta^g = -\beta_0 g^3/16\pi^2 - \beta_1 g^5/(16\pi^2)^2 - \cdots$ one has [8]

$$a = a_0 + \frac{n_V \beta_1}{8(16\pi^2)^3} g^4 + \mathcal{O}(g^6). \tag{1.1}$$

Here a_0 is the free field theory (one-loop) value of a and $n_V = \dim(\mathrm{Adj})$ is the number of vector fields. The result of LPR, that

$$\frac{da}{dt} \equiv -\mu \frac{da}{d\mu} = -c_g (\beta^g)^2, \qquad c_g = \frac{n_V}{32\pi^2 g^2}, \tag{1.2}$$

---

[1] More precisely "limit recursive flows." In what follows we refer to both limit cycles and limiting ergodic behavior simply as "cycles."

is incompatible with (1.1). We identify below the source of the error in the computation of LPR. We find that LPR neglect pertinent counterterms, and as a result they have in fact computed, to lowest order in perturbation theory, the rate of change not of a but of a different quantity, namely the function $\tilde{\beta}_b$ introduced by Jack and Osborn (JO) as a candidate for a c-function. Indeed, for single-coupling YM models JO have already shown that $\tilde{\beta}_b$ flows monotonically (to all orders in perturbation theory).

The quantity $\tilde{\beta}_b$ is related to a, which in JO is denoted by β_b, by

$$\tilde{\beta}_b = \beta_b + \tfrac{1}{8} w \beta^g .$$

Here w is a function of the coupling g, and $\beta^g = -dg/dt$ is the beta function. While $\tilde{\beta}_b$ and β_b agree on fixed points, the difference is parametrically large away from fixed points. In Section 2 we explain this in detail. There we use the method of LPR with care to include all necessary counterterms and rederive JO's consistency conditions, of which the monotonic flow of $\tilde{\beta}_b$ is but one example.

For models which display cycles the state of affairs is significantly more complex. In all these models the kinetic terms of the Lagrangian exhibit a "flavor" symmetry group G_F (that commutes with the gauge group). Scalar self-interactions and Yukawa couplings of scalars with fermions break G_F. The dependence of counterterms on the coupling constants characterizing these interactions is restricted by the pattern of breaking of G_F. There is a well-known, simple method of accounting for this. The coupling constants are treated as spurions, that is, as non-dynamical fields, and allowed to transform under G_F precisely so as to render the Lagrangian invariant under these symmetry transformations. Then, if the regulator respects the symmetry, so will the counterterms. It follows that the entries in the trace anomaly respect the symmetry too. As a is the coefficient of the G_F-invariant Euler density, it is itself G_F-invariant as well. And since the flow on a cycle corresponds to a G_F-transformation of the couplings, a remains constant on the cycle.

This begs the following question: "how is the monotonic flow of $\tilde{\beta}_b$ consistent with the constancy of a?" Actually, $\tilde{\beta}_b$ is also G_F-invariant, and is thus also guaranteed to be constant along the cyclic flow. The answer is found in the flow equation for $\tilde{\beta}_b$ given by Osborn in [9]. His equation is a generalization, applicable to these more complex theories, of that found in JO. This flow equation is not guaranteed to give monotonic flows, but can and does give constancy of $\tilde{\beta}_b$ along cycles. We review the work of Jack and Osborn concerning these more complex theories in Section 3, and show that $\tilde{\beta}_b$ remains constant on cycles. We remark there that while we agree with JO's mathematical formulae, we disagree with their interpretation of some of their results. In particular, JO give a monotonic flow, but not along an RG flow, which however they interpret as such.

As we have said, LPR presented an argument that gives monotonicity of the flow of a. They then argued that the infinite decrease of a on a cycle should be compensated by a counterterm, but that in its absence it must be that no four-dimensional theory may exhibit cycles. It is a delicious irony that it is precisely the neglect of counterterms in the treatment of LPR that leads them to the wrong conclusion that the flow of a (or even $\tilde{\beta}_b$) is monotonic.

In light of the lack of even a perturbative argument for the monotonic decrease of $\tilde{\beta}_b$ along RG flows, the results of CDGM and KS appear more remarkable. So in Section 5 we take on the task of extending the result of KS to flows between cycles or fixed points, as announced at the top of this introduction. We first use a non-RG flow on which a quantity $\widetilde{B}_b$ decreases monotonically, at least in perturbation theory. This quantity agrees with β_b on fixed points and cycles and we demonstrate that if an RG flow exists between fixed points/cycles, then the new non-RG flow also links those same fixed points/cycles. This non-RG flow captures the physics of scale-invariant theories. For example, all fixed points and cycles of the RG flow map to fixed points of the non-RG flow. The monotonic decrease of $\widetilde{B}_b$ gives still only a perturbative proof of a weak version of the generalized c-theorem. To go beyond perturbation theory we use the method of KS. The only additional assumption is that there exists a dimension-three current that is defined everywhere in theory space, that is, for any value of the coupling constants, such that it vanishes or is conserved at fixed points and generates flows on cycles.

The natural candidate for this current was introduced formally by JO. It corresponds to a particular element of the Lie algebra of G_F they call S. JO showed by direct computation that S vanishes to three loops if the field content of the QFT consists solely of scalars, and to two loops if of scalars and Dirac spinors. On the other hand, the element Q of the Lie algebra of G_F that generates the flow along the cycle is found in our computations to arise at three-loop order in gauge theories that include both scalars and spinors [6]. Could it be, then, that S is non-zero at three loops in these theories? And if so, what is the relation between Q and S? In Section 4 we take on the task of computing the lowest-order contribution to JO's S for the most general four-dimensional QFT, compare with Q and demonstrate that S is the interpolating function we need to complete the extension of the KS proof of a weak version of the c-theorem to cycles.

Let us specify here that we follow closely the notation of JO [8], with some notable exceptions. From here on, following JO, in the anomaly equation we use β_a and β_b for c and a respectively, although we still use the terminology c-theorem instead of the more accurate β_b-theorem. Also, we call $\lambda_{a,b,c}$ rather than a, b, c the infinite counterterms that give rise to $\beta_{a,b,c}$ (having infinite counterterms labeled by a and c can certainly produce confusion with the corresponding "beta functions" that appear in the Weyl anomaly and that are commonly referred to as a and c). Throughout this paper RG time is defined by $t = -\ln(\mu/\mu_0)$, so that it increases as we flow to the IR.

Many of our results are extracted from the work of JO. So it is perhaps necessary to remark that, besides parsing the results of JO to hopefully make them slightly more accessible to the general reader, we have made several novel contributions: we have discovered where in the argument of KS and LPR the quantity β_b is replaced by $\tilde{\beta}_b$ (or, in more generality, by $\widetilde{B}_b$); we have shown that the current in JO provides a generalization of the virial away from scale-invariant trajectories, and is conserved at fixed points; we have extended JO's calculation of S to third loop order, which is the leading non-vanishing contribution, in a Yang–Mills theory with scalars and spinors; and finally, and most remarkably, we have used the above to prove an extension of the weak version of the c-theorem, that $(\beta_b)_{\rm UV} > (\beta_b)_{\rm IR}$, for presumed RG flows that can now originate or terminate on cycles as well as fixed points, valid even outside perturbation theory (provided the implicit assumptions in KS do not invalidate their result).

2. Weyl consistency conditions

In this section we review the derivation of the Weyl consistency conditions of JO. The method uses as a starting point the expressions of Weyl invariance used by KS and by LPR. The presentation is formulated so that it becomes clear that the assumptions in those works already implicitly lead to the JO consistency conditions. Hence, although the derivation presented here may seem novel, it actually follows closely JO. We have included it here for completeness, for pedagogy and because it makes clear that neither the results of KS nor those of LPR should be in conflict with those of JO.

Let us briefly review Osborn's argument for the consistency conditions [9]. These are analogous to the well-known Wess–Zumino consistency conditions [10]. Let $\Delta_\sigma \widetilde{W}$ denote the action of a Weyl transformation on $\widetilde{W}$, the generating functional for connected renormalized Green functions. Because of the Abelian nature of the Weyl group, the Weyl consistency conditions follow:

$$[\Delta_\sigma, \Delta_{\sigma'}]\widetilde{W} = 0.$$

In JO the same consistency conditions are obtained by requiring finiteness of the trace of the stress-energy tensor in curved background and with spacetime-dependent couplings. One can also obtain the Weyl consistency conditions based on the arguments of LPR.

LPR start from a quantum action S_0 which is a function of a conformally flat metric, $\gamma_{\mu\nu} = e^{-2\tau(x)}\eta_{\mu\nu}$ and coupling constants $g^i(\mu)$ (in $d = 4-\epsilon$ regularization, with, say, minimal subtraction (MS)). By rescaling the fields, which are dummy variables of integration anyway, by $\phi \to (e^\tau)^\delta \phi$, where δ is the canonical dimension of the field (as in $\delta = (d-2)/2$ for scalars), and using the μ-independence of the bare couplings, one sees that the τ-dependence in S_0 arises only due to the scale dependence of renormalized coupling constants, $g^i(e^\tau \mu)$. Effectively, the regularized generating functional W satisfies

$$W[e^{-2\tau(x)}\eta_{\mu\nu}, g^i(\mu)] = W[\eta_{\mu\nu}, g^i(e^{\tau(x)}\mu)]. \tag{2.1}$$

Alternatively, Komargodski [11] argues that the functional is made invariant under Weyl transformations by adding a conformal compensator $\tau(x)$. One can write

$$W[e^{-2\tau(x)}\eta_{\mu\nu}, g^i(e^{-\tau(x)}\mu)] = W[\eta_{\mu\nu}, g^i(\mu)],$$

or, equivalently, that the left-hand side is invariant under $\tau \to \tau + \sigma$. Note that invariance is guaranteed by adding the conformal compensator and including in W all possible counterterms that are functions of spacetime-dependent background and coupling constants, $\gamma_{\mu\nu}(x)$ and $g^i(x)$. It is from counterterms that do not vanish for spacetime-independent coupling constants that the $\beta_{a,b,c}$-anomalies arise. It is convenient, in order to keep track of curvature-dependent terms, to do this in a more general background metric,

$$W[e^{-2\tau(x)}\gamma_{\mu\nu}(x), g^i(\mu)] = W[\gamma_{\mu\nu}(x), g^i(e^{\tau(x)}\mu)], \tag{2.2a}$$

$$W[e^{-2\tau(x)}\gamma_{\mu\nu}(x), g^i(e^{-\tau(x)}\mu)] = W[\gamma_{\mu\nu}(x), g^i(\mu)]. \tag{2.2b}$$

At the risk of restating the trivial, let us emphasize that it is not consistent to neglect the spacetime dependence of couplings when studying Weyl anomalies, since the Weyl transformation involves spacetime-dependent couplings. The counterterms associated with spacetime derivatives of these couplings will lead to additional anomalies. Some of these may—and as we will see, do—contribute to the dilaton/compensator scattering amplitude even after one takes the limit of flat background and spacetime-independent coupling constants.

The approach of LPR allows one to compute quantities associated with a model in a curved background with spacetime-independent coupling constants in terms of corresponding quantities for the same model but in a flat background with, however, spacetime-dependent coupling constants. This observation is not new. For example, in JO the same observation is used precisely for the same purpose, namely, to compute the anomalies associated with scale transformations using only computations in flat space. Similarly, the approach of Komargodski allows for an explicit nonlinear realization of scale invariance, at the price of introducing spacetime-dependent coupling constants. To reiterate, in either case it is important to realize that new counterterms are required to render the model finite, much like counterterms involving derivatives of the metric need to be introduced to render finite the model in a curved background. These new counterterms must involve derivatives of the coupling constants and lead to new Weyl anomalies. At the end of this section we study how these new anomalies contribute to the Wess–Zumino action for the conformal compensator $\tau(x)$ even after the couplings and the metric are taken to be spacetime-independent. For the remainder of this section we take a closer look at these counterterms, the anomalies they produce and the relations between them, that is, the JO consistency conditions, that follow from (2.2).

Consider the theory in the background of an arbitrary metric $\gamma_{\mu\nu}(x)$ and arbitrary spacetime-dependent coupling constants $g^i(x)$ corresponding to interaction terms $g^i(x)\mathcal{O}_i(x)$ in the Lagrangian. The arbitrary spacetime dependence of the couplings allows one to use them as sources for operators in the interaction part of the Lagrangian, by taking functional derivatives of the generating functional with respect to $g^i(x)$. Let $\widetilde{W}$ stand for the renormalized generating functional. It is convenient to separate the counterterms that are independent of quantum fields from the rest of the action. They can be taken out of the functional integral and contribute directly to $\widetilde{W}$:

$$\widetilde{W} = W + W_{\text{c.t.}}.$$

The generating functional W results from performing the functional integral over quantum fields in the absence of the quantum-field-independent counterterms. The counterterms required to render the theory finite were first classified in JO. They consist of all possible diff-invariant dimension-four operators constructed out of the metric and couplings and their derivatives:

$$W_{\text{c.t.}} = -\int \sqrt{-\gamma}\, \mu^{-\epsilon} \lambda \cdot \mathscr{R},$$

where dimensional regularization is used with $d = 4 - \epsilon$ and

$$\begin{aligned} \lambda \cdot \mathscr{R} = \lambda_a F + \lambda_b G + \lambda_c H^2 + \mathscr{E}_i \partial_\mu g^i \partial^\mu H + \tfrac{1}{2}\mathscr{F}_{ij} \partial_\mu g^i \partial^\mu g^j H + \tfrac{1}{2}\mathscr{G}_{ij} \partial_\mu g^i \partial_\nu g^j G^{\mu\nu} \\ + \tfrac{1}{2}\mathscr{A}_{ij} \nabla^2 g^i \nabla^2 g^j + \tfrac{1}{2}\mathscr{B}_{ijk} \partial_\mu g^i \partial^\mu g^j \nabla^2 g^k + \tfrac{1}{4}\mathscr{C}_{ijkl} \partial_\mu g^i \partial^\mu g^j \partial_\nu g^k \partial^\nu g^l. \end{aligned} \tag{2.3}$$

Here F is the Weyl tensor squared, G is the Euler density, H is proportional to the Ricci scalar, and $G_{\mu\nu}$ is the Einstein tensor:

$$\begin{aligned} F &= R^{\mu\nu\rho\sigma} R_{\mu\nu\rho\sigma} - \frac{4}{d-2} R^{\mu\nu} R_{\mu\nu} + \frac{2}{(d-2)(d-1)} R^2, \\ G &= \frac{2}{(d-3)(d-2)} (R^{\mu\nu\rho\sigma} R_{\mu\nu\rho\sigma} - 4 R^{\mu\nu} R_{\mu\nu} + R^2), \\ H &= \frac{1}{d-1} R, \qquad G_{\mu\nu} = \frac{2}{d-2}(R_{\mu\nu} - \tfrac{1}{2}\gamma_{\mu\nu} R). \end{aligned}$$

The quantities above are defined as in JO, with inessential d-dependent factors for later convenience. Each of the counterterms in $\lambda \cdot \mathscr{R}$ is an expansion in $1/\epsilon$ chosen to render $\widetilde{W}$ finite—for this one must in addition introduce wave-function and coupling constant counterterms, as usual. The coefficients λ are in general functions of the couplings $g^i(x)$.

The anomalous variation of the generating functional is dictated by these counterterms. While W satisfies (2.1) and (2.2), this is not true of $W_{\text{c.t.}}$, as can be seen by explicit computation. The anomaly is precisely the statement that the infinitesimal transformation $\tau \to \tau + \sigma$ in (2.2b),

$$\Delta_\sigma W_{\text{c.t.}} = W_{\text{c.t.}}[(1 - 2\sigma)\gamma_{\mu\nu}, g^i - \sigma\mu(dg^i/d\mu)] - W_{\text{c.t.}}[\gamma_{\mu\nu}, g^i],$$

fails to vanish. The anomalous variation can be split into a term that would occur even if σ were spacetime-independent plus a term proportional to the derivative of σ:

$$\Delta W_{\text{anomaly}} = \Delta_\sigma W_{\text{c.t.}} = -\int \sqrt{-\gamma}\, (\sigma \beta_\lambda \cdot \mathscr{R} + \partial_\mu \sigma\, \mathscr{Z}^\mu). \tag{2.4}$$

These terms again can be expanded using dimensional analysis and diff-invariance:

$$\begin{aligned}\beta_\lambda \cdot \mathscr{R} = \beta_a F + \beta_b G + \beta_c H^2 + \chi^e_i \partial_\mu g^i \partial^\mu H + \tfrac{1}{2}\chi^f_{ij}\partial_\mu g^i \partial^\mu g^j H + \tfrac{1}{2}\chi^g_{ij}\partial_\mu g^i \partial_\nu g^j G^{\mu\nu} \\ + \tfrac{1}{2}\chi^a_{ij}\nabla^2 g^i \nabla^2 g^j + \tfrac{1}{2}\chi^b_{ijk}\partial_\mu g^i \partial^\mu g^j \nabla^2 g^k + \tfrac{1}{4}\chi^c_{ijkl}\partial_\mu g^i \partial^\mu g^j \partial_\nu g^k \partial^\nu g^l,\end{aligned} \tag{2.5}$$

and[2]

$$\begin{aligned}\mathscr{Z}_\mu = G_{\mu\nu} w_i \partial^\nu g^i + \tfrac{1}{3}\partial_\mu(dR) + \tfrac{1}{3}R Y_i \partial_\mu g^i \\ + \partial_\mu(U_i \nabla^2 g^i + \tfrac{1}{2}V_{ij}\partial_\nu g^i \partial^\nu g^j) + S_{ij}\partial_\mu g^i \nabla^2 g^j + \tfrac{1}{2}T_{ijk}\partial_\nu g^i \partial^\nu g^j \partial_\mu g^k,\end{aligned} \tag{2.6}$$

up to terms with vanishing divergence. Since the anomaly gives a correction to a relation between finite Green functions (see 2.15), the anomalous terms (2.5) and (2.6) must themselves be finite.

Calculations of the coefficients in $\beta_\lambda \cdot \mathscr{R}$ and $\mathscr{Z}_\mu$ can be done using standard techniques of dimensional regularization with a mass-independent renormalization scheme, say MS. For now, let us concentrate on the relatively straightforward computation of $\beta_\lambda \cdot \mathscr{R}$. Since for constant σ the transformation $\delta\gamma_{\mu\nu} = -2\sigma\gamma_{\mu\nu}$ just counts dimensions, and the dimension of the volume element is d while that of the operators in $W_{\text{c.t.}}$ is four, we obtain

$$(\epsilon - \hat\beta^i \hat\partial_i)\lambda \cdot \mathscr{R} = \beta_\lambda \cdot \mathscr{R}, \tag{2.7}$$

where the beta function is

$$\mu \frac{dg^i}{d\mu} = \hat\beta^i = -\epsilon k^i g^i + \beta^i(g) \qquad \text{(no sum over } i\text{)}.$$

Here the derivative is taken holding the bare parameters fixed. k^i is defined by requiring that the Lagrangian scales appropriately: for $\phi' = \mu^{\delta\epsilon}\phi$ and $g'^i = \mu^{k^i\epsilon} g^i$, then $\mathscr{L}(\phi', g') = \mu^{-\epsilon}\mathscr{L}(\phi, g)$. Note that $\hat\beta^i\hat\partial_i \equiv \hat\beta^i \hat\partial/\hat\partial g^i$ denotes substitution of g^i by β^i wherever g^i may be found, e.g., $\hat\beta^i\hat\partial_i(\partial_\mu g^j) \equiv \partial_\mu \beta^j = \partial_i \beta^j \partial_\mu g^i$, and of course respects the standard rules of differentiation. Using (2.7), it is straightforward to show that, e.g.,

$$\chi^a_{ij} = (\epsilon - \hat\beta^k \partial_k)\mathscr{A}_{ij} - \mathscr{A}_{ik}\partial_j \hat\beta^k - \mathscr{A}_{jk}\partial_i \hat\beta^k.$$

The consistency conditions of JO can be understood as following from requiring that (2.2) applied to the complete renormalized generating function $\widetilde{W}$ fails only up to the finite, anomaly terms. The left-hand side of (2.2a) does not involve any counterterms from spacetime-dependent couplings, while the right-hand side does not involve any from a curved background. Hence, the counterterms in one and the other case must be related. Consider on the right-hand side of (2.2a), for example, the counterterm

$$\tfrac{1}{2}\mathscr{A}_{ij}\nabla^2 g^i \nabla^2 g^j = \tfrac{1}{2}\mathscr{A}_{ij}\hat\beta^i\hat\beta^j(\nabla^2\tau)^2 + \cdots, \tag{2.8}$$

---

[2] The second term involves the function of coupling constants d, which is not to be confused with $d = 4 - \epsilon$. We follow Osborn in this unfortunate choice of notation, hoping that with this warning no confusion will arise in what follows.

where we have expanded to lowest order in $\tau(x)$. Comparing with the counterterms on the left-hand side of (2.2a), that arise solely from a curved background, we have,

$$\sqrt{-\gamma}\,(\lambda_a F + \lambda_b G + \lambda_c H^2) = 8\lambda_b\left[(\nabla^2\tau)^2 - (\partial_\mu\partial_\nu\tau)^2 + \cdots\right] + 4\lambda_c\left[(\nabla^2\tau)^2 + \cdots\right]. \tag{2.9}$$

The λ_b term is a total derivative so for localized $\tau(x)$ it does not contribute (recall there is an implicit spacetime integration). Matching the terms in (2.8) and (2.9) we find that the counterterms are related,

$$4\lambda_c \sim \tfrac{1}{2}\mathscr{A}_{ij}\hat{\beta}^i\hat{\beta}^j, \tag{2.10}$$

where the symbol $\sim$ denotes equality up to finite terms, that is, the difference is finite as $\epsilon \to 0$. This precisely corresponds to Eq. (3.12) of JO. Applying $\mu\, d/d\mu$ on the bare couplings to derive RGEs and the corresponding beta functions, one then derives from this JO's consistency condition (3.21a),

$$8\beta_c = \chi^a_{ij}\beta^i\beta^j - \beta^i\partial_i X, \tag{2.11}$$

where X arises from the finite difference between the left- and right-hand sides of (2.10), and β_c and χ^a_{ij} are beta functions for λ_c and $\mathscr{A}_{ij}$, respectively. The remaining consistency conditions in JO can be obtained in a similar fashion. We only quote here one other consistency condition that plays an important role in what follows. Using (2.2a) the lowest order terms in $\tau(x)$ that are linear in the Einstein tensor give

$$8\partial_i\lambda_b \sim \mathscr{G}_{ij}\hat{\beta}^j. \tag{2.12}$$

If the finite difference is denoted by w_i, then one obtains

$$8\partial_i\beta_b = \chi^g_{ij}\beta^j - \beta^j\partial_j w_i - \partial_i\beta^j w_j.$$

This consistency condition is the origin of the proposal in JO for a c-function,

$$\tilde{\beta}_b \equiv \beta_b + \tfrac{1}{8}\beta^i w_i,$$

which satisfies

$$\partial_i\tilde{\beta}_b = \tfrac{1}{8}(\chi^g_{ij} + \partial_{[i}w_{j]})\beta^j, \tag{2.13}$$

where $\partial_{[i}w_{j]} = \partial_i w_j - \partial_j w_i$. Then its RG flow is monotonic provided the “metric” χ^g_{ij} is positive-definite, for

$$-\frac{d\tilde{\beta}_b}{dt} = \beta^i\partial_i\tilde{\beta}_b = \tfrac{1}{8}\chi^g_{ij}\beta^i\beta^j.$$

To summarize, the extension (2.2) of the invariance requirement of LPR in (2.1), when applied to *the complete set of counterterms required for finiteness when coupling constants have spacetime dependence*, leads to the consistency conditions of JO.

2.1. The trace anomaly and the computation of $\nabla_\mu \mathscr{Z}^\mu$

As formulated, the renormalized generating functional $\widetilde{W}$ is a function of the background metric and of renormalized spacetime-dependent coupling constants. As such we can obtain Green functions by functional differentiation,

$$T_{\mu\nu}(x) = \frac{2}{\sqrt{-\gamma}} \frac{\delta \widetilde{W}}{\delta \gamma^{\mu\nu}(x)} \qquad \text{and} \qquad \mathcal{O}_i(x) = \frac{1}{\sqrt{-\gamma}} \frac{\delta \widetilde{W}}{\delta g^i(x)} \ . \tag{2.14}$$

These definitions are to be understood as expressions for insertions in Green functions, so that, for example, $\mathcal{O}_i(x)$ stands for the renormalized insertion of the operator which may differ from the operator monomial in an expectation value (JO make this distinction explicit by referring to the Green function insertion as $[\mathcal{O}_i(x)]$, but we refrain from making that distinction to streamline notation knowing that there is little chance of confusion).

Using (2.14) in (2.2) and (2.4) one obtains

$$T^\mu_{\ \mu} = \hat{\beta}^i \mathcal{O}_i - \mu^{-\epsilon} \beta_\lambda \cdot \mathscr{R} + \mu^{-\epsilon} \nabla_\mu \mathscr{Z}^\mu. \tag{2.15}$$

This is the well-known trace anomaly, accounting for the effects of curved background and spacetime-dependent coupling constants. However, this equation is not quite correct in the most generality: there are two terms missing on the right-hand side. The first is an operator that vanishes by the equations of motion times the anomalous dimension of the corresponding quantum field. We have lost track of this term because the relation (2.1) is only correct up to terms that vanish by the equations of motion. The second missing term is more subtle: we have missed counterterms that may be needed to render some theories finite. When the kinetic terms of the Lagrangian exhibit a continuous symmetry the current associated with this symmetry is a dimension-three operator and a new type of counterterm is required in the presence of spacetime-dependent couplings, that is, a counterterm proportional to the product of the current and the derivative of a coupling. This will be discussed extensively, and the anomaly equation will be fixed, in Section 3.

Let us turn to the computation of $\mathscr{Z}^\mu$ in (2.4). It follows, of course, straightforwardly from the definition (2.4). Slightly less trivial is the fact that the computation must give a finite current $\mathscr{Z}^\mu$. That this must be so can be seen from the trace anomaly in (2.15), in which all other terms are already finite. This means that there must be cancellations among infinite terms that contribute to $\mathscr{Z}^\mu$. In fact, these cancellations are nothing but the consistency conditions, e.g., (2.10) and (2.12). For example, the terms in (2.4) involving the Einstein tensor (modulo terms that do not vanish for spacetime-independent σ) are

$$\begin{aligned}\int \sqrt{-\gamma}\, G^{\mu\nu}(-8\lambda_b \nabla_\mu \partial_\nu \sigma - \mathscr{G}_{ij} \partial_\mu g^i \hat{\beta}^j \partial_\nu \sigma) &= \int \sqrt{-\gamma}\, \partial_\nu \sigma\, G^{\mu\nu} \partial_\mu g^i (8 \partial_i \lambda_b - \mathscr{G}_{ij} \hat{\beta}^j) \\ &= \int \sqrt{-\gamma}\, \partial_\nu \sigma\, G^{\mu\nu} \partial_\mu g^i\, w_i.\end{aligned}$$

Here, in going from the first to the second line we used the finiteness condition (2.12) and the definition that the finite difference is w_i. Thus we have reproduced the first term in $\mathscr{Z}_\mu$ of (2.6). The remaining terms in (2.6) can be similarly found.

2.2. Wess–Zumino action

The Wess–Zumino action, W_{WZ}, is some function of $\tau(x)$ that will give $-\Delta W_{\mathrm{anomaly}}$ upon a Weyl transformation, $\tau(x) \to \tau(x) + \sigma(x)$. Focusing on the β_b-term in $\Delta W_{\mathrm{anomaly}}$,

$$-\int \sqrt{-\gamma}\, \sigma \beta_b G, \tag{2.16}$$

KS write the corresponding WZ term as[3]

$$\int \sqrt{-\gamma} \left\{ \tau \beta_b G - 4\beta_b \left[G^{\mu\nu} \tau_{,\mu} \tau_{,\nu} + \tau_{,\mu} \tau^{,\mu} \nabla^2 \tau + \tfrac{1}{2} (\tau_{,\mu} \tau^{,\mu})^2 \right] \right\}, \tag{2.17}$$

where we have introduced the shorthand $\tau_{,\mu} = \partial_\mu \tau$. However, this computation is incomplete. The problem with this is that we have ignored the effect of the new counterterms arising from spacetime dependence of the couplings. Since we will not need a Wess–Zumino action for our generalization of the KS argument to theories with cycles, we will not aim at being complete and only point out one interesting consequence here. Consider, for example, the term in $\mathscr{Z}^\mu$

$$-\int \sqrt{-\gamma}\, \partial_\mu \sigma \, w_i G^{\mu\nu} \partial_\nu g^i. \tag{2.18}$$

Now with $\partial_\mu g^i = -\beta^i \tau_{,\mu}$, one has the following generalization of the Wess–Zumino dilaton action:

$$\int \sqrt{-\gamma} \left\{ \beta_b \tau G - 4(\beta_b + \tfrac{1}{8} w_i \beta^i) \left[G^{\mu\nu} \tau_{,\mu} \tau_{,\nu} + \tau_{,\mu} \tau^{,\mu} \nabla^2 \tau + \tfrac{1}{2} (\tau_{,\mu} \tau^{,\mu})^2 \right] \right\}. \tag{2.19}$$

The Weyl variation of (2.19) gives the sum of (2.16) and (2.18) (if $\partial_\mu g^i = -\beta^i \tau_{,\mu}$ there). The correction that takes β_b into $\tilde{\beta}_b = \beta_b + \frac{1}{8} w_i \beta^i$ is generally of lower order than the running of β_b. That is, $\frac{1}{8} w_i \beta^i$ is of lower order than $\beta_b - \beta_{b0}$, where β_{b0} stands for the free field theory value of β_b.

Let us be more explicit. Consider the perturbative result of JO for a YM theory with gauge coupling g,

$$\beta_b = \beta_{b0} + \frac{n_V \beta_1}{8(16\pi^2)^3} g^4 + \mathcal{O}(g^6). \tag{2.20}$$

JO also give $gw = 2n_V/16\pi^2 + \cdots$, and therefore

$$\tilde{\beta}_b = \beta_{b0} - \frac{n_V \beta_0}{4(16\pi^2)^2} g^2 + \mathcal{O}(g^4).$$

---

[3] The sign in the term cubic in τ is opposite to that of KS because we use the opposite sign convention for the conformal compensator, which gives Δ_σ as in JO.

Notice however that, as already mentioned in the introduction, LPR give

$$\frac{d\beta_b}{dt} \overset{\text{LPR}}{=} -\frac{n_V}{32\pi^2 g^2}(\beta^g)^2,$$

which is incompatible with (2.20). However, LPR would have obtained the correct result, namely that they computed in fact the rate of change of $\tilde{\beta}_b$, had they realized the dilaton scattering amplitude is given by the improved Wess–Zumino action in (2.19).

It is instructive to compare the one other element of the LPR computation with the corresponding one in JO. The coefficient c_g of LPR appears to play the role of the "metric" χ^g_{ij} in the consistency condition (2.13). In Appendix A we show that c_g is actually related to JO's $-2\chi^a_{ij}$. We have already encountered χ^a_{ij} as a metric in (2.11). Explicit computations show that $-2\chi^a_{ij}$ agrees with χ^g_{ij} to second order in perturbation theory in any four-dimensional theory—see Appendix A. Since χ^a_{ij} is negative-definite, were $\chi^g_{ij} = -2\chi^a_{ij}$ to hold to all orders in perturbation theory it would ensure positivity of χ^g_{ij} to all orders. However, this is not the case: as we show in Appendix A the equality fails, for example, at third order in a YM theory with a single gauge coupling.

3. Flavor symmetries, dimension-three operators and the corrected trace anomaly

As we have mentioned above, in deriving the trace anomaly we have missed counterterms that may be needed to render some theories finite. When the kinetic terms of the Lagrangian exhibit a continuous symmetry the current associated with this symmetry is a dimension-three operator and a new type of counterterm is required in the presence of spacetime-dependent couplings, that is, a counterterm proportional to the product of the current and the derivative of a coupling.

Consider a theory with n_S real scalar fields. The kinetic part of the Lagrangian

$$\mathscr{L}_K = \tfrac{1}{2}\partial^\mu\phi_a\partial_\mu\phi_a$$

exhibits a continuous symmetry under transformations of the fields $\delta\phi_a = \omega_{ab}\phi_b$, where ω is in the algebra of the flavor group $G_F = SO(n_S)$. In order to renormalize this theory in the presence of spacetime-dependent coupling constants one must introduce a new counterterm of the form

$$\mathscr{L}_{\text{c.t.}} = (\partial^\mu g^i)(N_i)_{ab}\phi_b\partial_\mu\phi_a, \tag{3.1}$$

with $(N_i)_{ab} = -(N_i)_{ba}$, that is, in the algebra of G_F. Note that this new counterterm is not accounted for in $W_{\text{c.t.}}$ which by construction is independent of quantum fields. Note also that additional counterterms, symmetric under $a \leftrightarrow b$, must also be introduced. One may integrate by parts to write these as terms with no derivatives acting on the quantum fields. While necessary, they do not play a central role in what follows.

To be more explicit, we consider a theory of real scalars and write for the bare Lagrangian

$$\mathscr{L} = \tfrac{1}{2}\gamma^{\mu\nu} D_\mu\phi_{0a} D_\nu\phi_{0a} + \tfrac{1}{8}(d-2)\phi_{0a}\phi_{0a}H - \tfrac{1}{4!}g^0_{abcd}\phi_{0a}\phi_{0b}\phi_{0c}\phi_{0d}. \tag{3.2}$$

This is written in term of bare fields ϕ_0. The second term is introduced to ensure conformal invariance of the classical action. In the potential term, the bare couplings g^0_{abcd} are completely symmetric under exchange of the indices a, b, c and d. The covariant derivative,

$$D_\mu\phi = (\partial_\mu + A_{0\mu})\phi,$$

is introduced with an eye towards including the counterterm (3.1),

$$A_{0\mu} = A_\mu + (D_\mu g)_I N_I. \tag{3.3}$$

Here, following JO, we use the compact notation $I = (abcd)$ and we have left implicit the Lie-algebra indices (so that $N_I^T = -N_I$ and $A_\mu^T = -A_\mu$). Note that N_I is a function of the renormalized couplings that has an ϵ-expansion starting at order $1/\epsilon$. If the theory contains gauge fields and some of the scalars are charged under the gauge group $G_g \subseteq G_F$, it is straightforward to include an additional quantum gauge field in addition to the background field A_μ.

The Lagrangian (3.2) is explicitly locally G_F-symmetric if we agree to transform the couplings and the gauge fields:

$$\begin{aligned} \delta g_{abcd} &= \omega_{ae} g_{ebcd} + \text{permutations} \qquad (\delta g_I = (\omega g)_I \text{ for short}),\\ \delta A_\mu &= D_\mu \omega. \end{aligned}$$

The first of these is already used in defining the covariant derivative $(D_\mu g)_I$ in (3.3). It is very important to note at this point that if this explicit local invariance is non-anomalous it can (and will) be used to constrain the counterterms and the generating functional $\widetilde{W}$,

$$\widetilde{W}[\gamma^{\mu\nu}(x), (\Omega g)_I(x), \Omega D_\mu \Omega^{-1}] = \widetilde{W}[\gamma^{\mu\nu}(x), g_I(x), A_\mu], \tag{3.4}$$

where $\Omega(x) = \exp(\omega(x)) \in G_F$. Of course, in theories without spinors the symmetry is trivially non-anomalous. Furthermore, derivatives of the generating functional with respect to the background field now give insertions of the scalar current.

It is not our intention to repeat the calculations of JO in their entirety here. We will instead describe the main ingredients and results. We have already described the two main new ingredients, namely, the need for new counterterms and the introduction of a background field to ensure invariance under G_F in (3.4). As before, additional quantum-field-independent counterterms are required. These are as in (2.3) but with the replacement $\partial_\mu \to D_\mu$ to ensure G_F invariance. Additional counterterms involving the field strength $F_{\mu\nu} = [D_\mu, D_\nu]$ are also required,

$$\tilde{\lambda}\cdot\mathscr{R} = \lambda\cdot\mathscr{R} + \tfrac{1}{4}\mathrm{Tr}(\mathscr{K} F_{\mu\nu}F^{\mu\nu}) + \tfrac{1}{2}\mathrm{Tr}(\mathscr{P}_{IJ}F^{\mu\nu})(D_\mu g)_I (D_\mu g)_J. \tag{3.5}$$

Moreover, as advertised, new field-dependent counterterms must be included,

$$\mathscr{Q} = \eta_{ab}\phi_a\phi_b H + (\delta_I)_{ab}\phi_a\phi_b(D^2 g)_I + \tfrac{1}{2}(\epsilon_{IJ})\phi_a\phi_b(D_\mu g)_I(D_\mu g)_J. \tag{3.6}$$

Proceeding much as before, JO find

$$T^\mu_{\ \mu} = \widehat{B}_I\mathcal{O}_I + \beta^{\mathscr{Q}} + (D^\mu\phi)^T B^A_\mu\phi - \mu^{-\epsilon}\beta_{\tilde{\lambda}}\cdot\mathscr{R} + \nabla_\mu(\widetilde{\mathscr{Z}^\mu} + J^\mu + J^\mu_\Theta) - ((1+\gamma+S)\phi)\cdot\frac{\delta}{\delta\phi}S_0.$$

Many comments are in order. The last term, involving the derivative of the full action integral S_0, vanishes by the equations of motion. We have included it here for completeness. We have already commented that a similar term is missing from (2.15). The operator $\mathcal{O}_I = -\frac{1}{4!}\phi_a\phi_b\phi_c\phi_d$ is precisely the interaction term in the Lagrangian, and its coefficient is defined as

$$\widehat{B}_I \equiv \hat{\beta}_I - (Sg)_I,$$

where $\hat{\beta}_I = \mu\, dg_I/d\mu$. The Lie-algebra element S is defined through

$$S = N_I\widehat{B}_I \qquad \text{or} \qquad S = -N^1_I g_I, \tag{3.7}$$

where N^1_I is the residue of the simple ϵ-pole in N_I. Similarly,

$$B^A_\mu \equiv \beta^A_\mu + D_\mu S \equiv \rho_I(D_\mu g)_I + D_\mu S \equiv P_I(D_\mu g)_I,$$

where the beta function for the background field A_ν is $\beta^A_\nu \equiv \mu\, dA_\nu/d\mu$. The beta functions for the field-dependent quadratic counterterms are

$$\beta^{\mathscr{Q}} \equiv \beta^\eta_{ab}\phi_a\phi_b H + (\beta^\delta_I)_{ab}\phi_a\phi_b(D^2 g)_I + \tfrac{1}{2}(\beta^\epsilon_{IJ})\phi_a\phi_b(D_\mu g)_I(D_\mu g)_J\,.$$

The term $\beta_{\tilde{\lambda}}\cdot\mathscr{R}$ is the obvious generalization of (2.5) while the current $\widetilde{\mathscr{Z}^\mu}$ is defined as in (2.6) but rendered G_F-invariant by replacing derivatives by covariant derivatives. In addition, $\widetilde{\mathscr{Z}^\mu}$ has contributions from the new counterterms in (3.5) and there are additional contributions to the terms with $\mathscr{A}$ and $\mathscr{B}$ that involve $\widehat{B}^A_\mu$. Finally, the current J^μ is

$$J^\mu = (D^\mu\phi)^T S\phi,$$

while the current J^μ_Θ arises from the counterterms in (3.6) and has a complicated expression in terms of the simple ϵ-poles in δ_I and ϵ_{IJ} (see JO for details [8, Eqs. (6.21–22)]).

For the special case of vanishing background field, spacetime-independent couplings and flat background the only terms that remain in the trace anomaly equation are

$$T^\mu_{\ \mu} - \partial_\mu J^\mu = \widehat{B}_I\mathcal{O}_I - ((1+\gamma+S)\phi)\cdot\frac{\delta}{\delta\phi}S_0. \tag{3.8}$$

We have written the divergence of the current J^μ on the left-hand side of the equation to highlight the similarity of this equation with that obtained when looking for scale but not conformally invariant theories. In that case one considers the dilatation current

$$\mathcal{D}^\mu = x_\nu T^{\mu\nu} - V^\mu$$

with the virial current given by

$$V^\mu = (D^\mu\phi)^T Q\phi,$$

so that

$$\partial_\mu \mathcal{D}^\mu = T^\mu_{\ \mu} - \partial_\mu V^\mu = (\hat{\beta}_I - (Qg)_I)\mathcal{O}_I - ((1+\gamma+Q)\phi)\cdot\frac{\delta}{\delta\phi}S_0.$$

In this case Q is determined by the condition that the divergence of the dilatation current vanishes. This only occurs for special values of Q and the coupling constants, namely those which solve

$$\hat{\beta}_I - (Qg)_I = 0.$$

It is clear that (3.8) suggests a relation between J^μ and V^μ. This relation can be easily understood from the behavior of QFT in curved spacetime. Indeed, it is well-known that Weyl-invariant theories in curved spacetime are conformal in the flat space limit [12]. The obstruction to lifting non-conformal scale-invariant theories to curved spacetime comes precisely from the existence of a non-trivial virial current. To account for the virial current, it is thus natural to introduce a background gauge field $A_\mu(x)$ which then allows for the general dilatation current

$$\mathcal{D}^\mu = x_\nu T^{\mu\nu} - J^\mu. \tag{3.9}$$

It is instructive to understand the origin of the current J^μ in JO's trace anomaly equation (3.8). The Weyl transformation for the generating functional of (2.2) now must account for the transformation of the background vector field A_μ (or, alternatively, for the presence of the new counterterm N_I):

$$W[e^{-2\tau(x)}\gamma_{\mu\nu}, A_\lambda(\mu), g_I(\mu)] = W[\gamma_{\mu\nu}, A_\lambda(e^{\tau(x)}\mu), g_I(e^{\tau(x)}\mu)], \tag{3.10a}$$

$$W[e^{-2\tau(x)}\gamma_{\mu\nu}, A_\lambda(e^{-\tau(x)}\mu), g_I(e^{-\tau(x)}\mu)] = W[\gamma_{\mu\nu}, A_\lambda(\mu), g_I(\mu)]. \tag{3.10b}$$

Focusing on, e.g., (3.10b), we have indicated that all of the renormalization-point dependence on the left-hand side, which enters also through the counterterm in A_λ, carries the conformal compensator factor. Concentrating on the new contribution, the variation under $\tau \to \tau + \sigma$ gives

$$\begin{aligned}\Delta_\sigma[\tfrac{1}{2}(D_\mu\phi)^T D^\mu\phi] &= -(D_\mu\phi)^T(N_I\hat{\beta}_I\partial^\mu\sigma)\phi + \cdots\\ &= -\partial_\mu\sigma\, J^\mu + \cdots\,,\end{aligned}$$

where the ellipses stand for the variations with σ that are accounted for elsewhere, and we have used in the last step the second equation (3.7). As we see, the natural outcome from a scaling transformation is not the canonical dilatation current $x_\nu T^{\mu\nu}$ but the current (3.9).

It is now obvious, as will be explicitly shown in the next section, that J^μ corresponds to the virial current V^μ on cycles. More precisely, J^μ is the appropriate virial current everywhere in parameter space. The natural dilatation current is thus defined by (3.9) and differs from the canonical one as soon as J^μ is non-trivial.

The divergence of the current J^μ in (3.8) precisely cancels the corresponding terms on the right-hand side, so that one has,

$$T^\mu_{\ \mu} = \hat{\beta}_I \mathcal{O}_I - ((1+\gamma)\phi) \cdot \frac{\delta}{\delta\phi} S_0, \tag{3.11}$$

which is nothing but the well-known expression for the trace anomaly in flat space. This is as expected because (2.2) are correct. However, in order to derive the consistency conditions in the most general case, one must use (3.10) instead of (2.2) to account for all possible counterterms. It is therefore not surprising that the work of KS led LPR to the conclusion that non-conformal scale-invariant theories are impossible: by neglecting these important counterterms both of them effectively assume that scale implies conformal invariance as a starting point. Let us turn anew to the JO consistency conditions.

In the general case considered here the JO consistency conditions are modified relative to what has been presented in Section 2. On the one hand the conditions have to be covariant under transformations by the symmetry group G_F. On the other, there are additional terms that arise from the additional counterterms required to render the theory finite. Osborn gives the form of these most general consistency conditions [9]. Two conditions play a role in our discussion:

$$\begin{aligned} 8\partial_I \beta_b &= \chi^g_{IJ} B_J - B_M \partial_M w_I - (\partial_I B_M) w_M - (P_I g)_M w_M \\ &= \chi^g_{IJ} B_J - \beta_M \partial_M w_I - (\partial_I \beta_M) w_M - (\rho_I g)_M w_M, \end{aligned} \tag{3.12}$$

and

$$B_I P_I = 0. \tag{3.13}$$

In addition, covariance under G_F gives, e.g.,

$$(\omega g)_I \partial_I \beta_b = 0 \qquad \text{and} \qquad (\omega g)_I \partial_I S = [\omega, S]. \tag{3.14}$$

Of course, the first of these applies to any G_F-invariant while the second to any antisymmetric tensor (for example, any Lie-algebra valued function). Using the first of (3.14) in (3.12) gives a nontrivial relation among several beta functions:

$$(\omega g)_I \left[\chi^g_{IJ} B_J - B_M \partial_M w_I - (\partial_I B_M) w_M - (P_I g)_M w_M\right] = 0, \tag{3.15}$$

or, equivalently,

$$(\omega g)_I \left[\chi^g_{IJ} B_J - \beta_M \partial_M w_I - (\partial_I \beta_M) w_M - (\rho_I g)_M w_M\right] = 0. \tag{3.16}$$

These conditions can be used to understand aspects of the flow of β_b. Consider the flow defined by some arbitrary function $f_I(g)$,

$$\frac{d\bar{g}_I}{d\eta} = -f_I(\bar{g}(\eta)).$$

If one takes $f_I = \beta_I$ then the flow can be identified with the RG flow, with $\eta = t = -\ln(\mu/\mu_0)$. From (3.12) we have

$$-8\frac{d\widetilde{B}_b}{d\eta} = \chi^g_{IJ} f_I B_J - f_I B_M \partial_{[M} w_{I]} - (P_I g)_M f_I w_M$$

where $\widetilde{B}_b = \beta_b + \frac{1}{8} B_I w_I$, and

$$-8\frac{d\tilde{\beta}_b}{d\eta} = \chi^g_{IJ} f_I B_J - f_I \beta_M \partial_{[M} w_{I]} - (\rho_I g)_M f_I w_M.$$

Three special cases are of most interest. Consider first $f_I(g) = (\omega g)_I$. From the second equation in (3.14) we see that on the flow S is constant. This is a recursive flow (cycle or ergodic). It follows from the G_F-invariance of $\widetilde{B}_b$ and $\tilde{\beta}_b$ that these remain constant on the flow. This is a consequence of the detailed cancellations that must be satisfied by the beta functions in (3.15) and (3.16). This general result can be applied to actual limit cycles, for which $\omega = Q$. Based on an incorrect equation, roughly (3.12) with S, w_I and ρ_I set to zero, LPR argue that cycles cannot occur in four-dimensional theories. They argue that β_b decreases monotonically, $d\beta_b/dt < 0$ on the cycle, which is inconsistent since there is no counterterm to remove the infinity as β_b flows to the IR. We now see that indeed there are counterterms, missed by LPR but known to JO, that not only account for the would-be divergence but in fact ensure G_F-covariance and hence constancy of β_b (and $\tilde{\beta}_b$) on the cycle.

The second special flow we consider is that of $f_I = \beta_I$, that is, actual RG-flow. Now we have

$$-8\frac{d\tilde{\beta}_b}{dt} = \chi^g_{IJ} \beta_I B_J - (\rho_I g)_M \beta_I w_M. \tag{3.17}$$

If we were to ignore the last term on the right-hand side we could establish a perturbative c-theorem for $\tilde{\beta}_b$. Indeed, to two loops $B_I = \beta_I$ and $\chi^g_{IJ} = -2\chi^a_{IJ} > 0$ so the right-hand side of (3.17) would be positive-definite along a perturbative flow. However, the last term is parametrically of the same order as the first on the right-hand side of (3.17) precluding a perturbative proof of the strong version of the c-theorem.

The third and last special case we wish to consider is that of $f_I = B_I$. In that case,

$$-8\frac{d\widetilde{B}_b}{d\eta} = \chi^g_{IJ} B_I B_J, \tag{3.18}$$

where we have used the consistency condition (3.13). This shows that there are flows on which some quantities decrease monotonically. However, these flows are not RG flows, even though they are perturbatively very close to RG flows. If one could establish that these flows have as endpoints the same fixed points (or limit cycles) as RG flows one would have a proof of the weak version of the c-theorem. We return to this question in Section 5.

It is worth noting that J^μ, although not conserved, has no anomalous dimension and thus dimension exactly three on a cycle. This observation is important because it contradicts the view, espoused by Osborn, that one can take B_I to be the beta function. If this were the case one would have conformal invariance on cycles (which would then be interpreted as fixed points), which however is impossible in light of unitarity and the existence of the non-vanishing non-conserved current J^μ of dimension three. It is clear that the relation between β_I and B_I is not simply a change of renormalization scheme, since some fixed points of B_I are not fixed points of β_I. For this reason, in the rest of this work we will continue to make explicit the distinction between RG-flows and η-flows generated by B_I.

4. S is Q (on cycles)

We alluded above to the similar role played by the current J^μ that arises in the trace anomaly and the virial current V^μ that defines a conserved dilatation current on a cycle. In this section we investigate the relation between them, or more specifically, the relation between the Lie-algebra valued function S and the Lie-algebra constant Q defined only on the limit cycle.

What is known about S? JO have shown, by direct calculation, and we have verified, that in a scalar field theory S vanishes up to third order in the loop expansion. The result holds even if gauge fields are included and the scalars are charged under the gauge group. For theories with scalars and fermions, JO have shown, and we have verified, that S remains zero to two loops. However, this is consistent with a possible equality of S and Q on cycles. Indeed, we have obtained previously that Q is of third order in the loop expansion in Yang–Mills theories with scalars and fermions, while in purely scalar theories we were unable to find a non-vanishing Q up to third order in the loop expansion.

As expected from the discussion above, we will show that (up to conserved current)

1. S is Q on cycles,
2. S vanishes at fixed points.

In light of these results the computation of Q can be tremendously simplified given an explicit expression for S. The procedure to determine Q involves determining first the beta functions for the coupling constants to second order in the loop expansion for scalar self-couplings, to third order in the loop expansion for Yukawa couplings and to fourth order in the loop expansion for Yang–Mills couplings, and then solving the system of nonlinear coupled equations $\beta^g = 0$ and $\beta_I = (Qg)_I$ (we implicitly use here that g_I can also stand for Yukawa couplings). Since S must have a perturbative expansion that starts at third order in the loop expansion, to determine Q from S it suffices to evaluate it with coupling constants on the cycle computed to lowest order in the loop expansion. So Q is obtained from S by determining the zeroes of the one-loop beta functions (two-loop for gauge couplings): if $S = 0$ on the zero of the beta functions, the zero is a fixed point of the RGE, but if $S \neq 0$ on the zero, then the zero is a point on a cycle and $Q = S$ there.

To this end an explicit, three-loop expression for S is required. But as pointed out above, there has been no computation of S to the order where one would expect it to be non-vanishing if S were to equal Q on cycles. We have endeavored to compute S to third order in the loop expansion for a general theory containing n_S real scalars and n_f Weyl spinors, possibly charged under a gauge group. The potential in the Lagrangian is

$$-V = \tfrac{1}{4!}\lambda_{abcd}\phi_a\phi_b\phi_c\phi_d + (\tfrac{1}{2}y_{a|ij}\phi_a\psi_i\psi_j + \text{h.c.}).$$

The details of the computation are spelled out in Appendix B. The surprisingly simple result is

$$(16\pi^2)^3 S_{ab} = \tfrac{5}{8}\operatorname{tr}(y_a y_c^* y_d y_e^*)\lambda_{bcde} + \tfrac{3}{8}\operatorname{tr}(y_a y_c^* y_d y_d^* y_b y_c^*) - \{a \leftrightarrow b\} + \text{h.c.}.$$

We have evaluated this expression on the fixed points and cycles of the theories we explored in [3, 5, 6] and found that in each case S vanishes at all fixed points and equals our previous determination of Q on all cycles. It should be mentioned that the models in [3,5] are in $d = 4 - \epsilon$ dimensions, with ϵ small but positive. In these cases the consistency equations still apply provided one carefully tracks the presence of ϵ in the beta functions, particularly in $\hat{\beta}^i = -\epsilon k^i g^i + \beta^i$ and correspondingly in $\widehat{B}^i$.

Now for the proof of the propositions above. First we show that $S = Q$ on cycles. Consider the η flow with $f_I = B_I$, with boundary condition that at $\eta = 0$ the point $\bar{g}_I(0)$ is on the cycle. Then $B_I(0) = ([Q - S]g(0))_I$ is in the Lie algebra of G_F and the left-hand side of (3.18) vanishes. Since $\chi^g_{IJ} = -2\chi^a_{IJ}$ to second order in the loop expansion, and $-2\chi^a_{IJ}$ is positive-definite, (3.18) gives $B_I(0) = 0$. This implies $S = Q + \Delta Q$ on cycles, where $(\Delta Q\, g)_I = 0$. But if $\Delta Q \neq 0$ this corresponds to a conserved current, and we are free to redefine the scale current by a conserved current, $Q \to Q + \Delta Q$. Hence, $S = Q$ on cycles.

For theories with two scalars there is an alternative, perhaps simpler proof that S equals Q when evaluated on a cycle. Consider (3.18) evaluated on a point on the cycle. It is easy to show that S is a constant on the cycle: $-dS/dt = \beta_I \partial_I S = (Qg)_I \partial_I S = [Q, S] = 0$, where the last two steps follow from (3.14) and the fact that, for two flavors, the flavor group, $SO(2)$, is Abelian. Now, as before, we consider the η-flow defined by the B_I function starting from a point on the RG-cycle (we make the distinction of the actual RG-cycle and a η-cycle explicit, to avoid confusion). The flow is defined by $-d\bar{g}/d\eta = B_I = \beta_I - (Sg)_I = ([Q - S]g)_I$, where the last step follows from assuming the initial point is on the RG-cycle and then noting that the solution corresponds to a trajectory that traverses the same cycle but at a different angular speed (the angular speeds are Q_{12} and $Q_{12} - S_{12}$ for the RG- and η-cycles, respectively). Therefore the η-cycle is generated by a trajectory in G_F and it follows that, just as for an RG-cycle, any G_F-invariant remains constant on the η-cycle. But the consistency condition (3.18) then implies that $B_I = ([Q-S]g)_I = 0$ on the cycle. Since Q and S are each characterized by a single number the only solution is $S_{12} = Q_{12}$ (on the cycle).

It is easy to show that $(Sg)_I = 0$ at a fixed point, and this is consistent with the notion that a fixed point corresponds to the case $(Qg)_I = 0$. To see this, notice that at a fixed point $B_I = -(Sg)_I$ so at that point the flow corresponds to a first order G_F-transformation. That is, the first derivative with respect to η of G_F-invariants vanishes at the fixed point. Hence, (3.18) gives that $\chi^g_{IJ}(Sg)_I(Sg)_J = 0$ and hence $(Sg)_I = 0$ at the fixed point. The solution is that either $S = 0$ at the fixed point, or there is an emergent symmetry at the fixed point, and J^μ is the corresponding conserved current. This completes the proof of the two propositions above.

5. An improved *c*-theorem

As we have seen, the consistency relations of JO lead to a monotonic η-flow in perturbation theory,

$$-\frac{d\widetilde{B}_b}{d\eta} = B_I \partial_I \widetilde{B}_b = \tfrac{1}{8}\chi^g_{IJ} B_I B_J \geq 0,$$

where we have defined $\widetilde{B}_b = \beta_b + \frac{1}{8} B_I w_I$. Only the last step in this sequence of relations invokes perturbation theory, for the positivity of the metric χ^g_{ij} is established perturbatively; see Appendix A. However, generally this η-flow does not correspond to an RG-flow, so one cannot conclude from this that the strong form of the c-theorem—that $c = \widetilde{B}_b$ decreases monotonically along the RG-flow—applies. One could try to use instead the consistency condition for the RG-flow of $\tilde{\beta}_b$ in (3.17), but the right-hand side of that condition is not necessarily positive-definite. Alternatively we use the fact that $B_I = 0$ at fixed points and on cycles, and show that there are η-flows between fixed points and cycles (or between fixed points and fixed points or between cycles and cycles) and hence prove the weak version of the c-theorem, that is, that $(\beta_b)_{\rm UV} > (\beta_b)_{\rm IR}$, where the subscript refers to β_b evaluated at the corresponding end-point (or end-limit-cycle) of the RG trajectory.

We assume there is an RG flow with endpoints on fixed points or limit cycles,

$$-\frac{dg_I}{dt} = \beta_I(g(t))$$

with $g_I(-\infty) = (g_I)_{\rm UV}$ and $g_I(\infty) = (g_I)_{\rm IR}$. In the case of limit cycles the end points refer rather to limit trajectories. Now, consider the solution to the η flow,

$$-\frac{d\bar{g}_I}{d\eta} = B_I(\bar{g}(\eta)).$$

It is given by

$$\bar{g}(\eta) = F(\eta) g(\eta),$$

where

$$F(\eta) = T\left(\exp\left[-\int_{-\infty}^{\eta} d\eta' S(\eta')\right]\right)$$

is an element of G_F and T is the η-ordered product. As such, $\beta_I(\bar{g}) = \beta_I(Fg) = (F\beta)_I(g)$ so that the fixed points and cycles $(g_I)_{\rm UV}$ and $(g_I)_{\rm IR}$ are mapped to physically equivalent fixed points and cycles $(\bar{g}_I)_{\rm UV}$ and $(\bar{g}_I)_{\rm IR}$. Since $\widetilde{B}_b$ is G_F-invariant and $B_I = 0$ on fixed points and cycles, it follows from the monotonic flow of $\widetilde{B}_b$ that $(\beta_b)_{\rm UV} > (\beta_b)_{\rm IR}$. We emphasize that this proof relies on perturbation theory to establish the positivity of the metric χ^g_{IJ}. For a non-perturbative proof we turn to the method of KS.

Let us review the argument of KS. Our presentation is closer in spirit to that of LPR. We will try to note explicitly when implicit assumptions in that argument are made. While plausible, these assumptions should be justified for the theorem to be established. We deviate from both presentations in that we do not derive nor use a Wess–Zumino dilaton action for, as we will see, this is not necessary for the computation.

Consider the four point function of the operator $\frac{1}{2}\partial_\mu\mathcal{D}^\mu = \frac{1}{2}\partial_\mu(x_\nu T^{\mu\nu} - J^\mu)$ in an arbitrary four-dimensional theory which is classically scale-invariant. In the absence of dimension-three currents this is just the four point function of $\frac{1}{2}T^\mu_{\ \mu}$, but we can more generally consider in one swoop the case where dimension-three currents are present. Furthermore, we will consider kinematics such that $p_i^2 = 0$, $i = 1, \ldots, 4$, for the momenta p_i of the four insertions. This is equivalent to computing the $\tau(x)$ scattering amplitude $\mathcal{A}(s,t)$ with the on-shell condition $\nabla^2\tau = 0$, so that the Mandelstam variables satisfy $s + t + u = 0$.

Now, we will assume that the forward scattering amplitude $\mathcal{A}_{\rm fwd}(s) = (s,0)$ exists, that is that the limit $t \to 0$ of (s,t) exists. This could fail if $\mathcal{A}(s,t)$ had terms of the form, e.g., $\sim s^2\ln(t)$. As we have explained, the response of the theory to Weyl rescaling is measured in the general case by $\partial_\mu\mathcal{D}^\mu$, so $\mathcal{A}_{\rm fwd}(s)$ can be computed by taking four $\tau(x)$-derivatives of the generating functional and then taking the metric as flat, the coupling constants to be spacetime-independent and the background field A_μ and the conformal compensator to vanish. Alternatively, and more straightforwardly, one can work with a conformally flat metric and having the only spacetime dependence in couplings and A_μ arise through the dependence on the conformal factor $\tau(x)$, so that one merely needs to take $\tau(x) = 0$ after four times differentiating $\widetilde{W}$. Now the first derivative simply gives the dilatation anomaly equation

$$\nabla_\mu\mathcal{D}^\mu = \widehat{B}_I\mathcal{O}_I + \beta^{\mathscr{Q}} + (D^\mu\phi)^T B^A_\mu\phi - \mu^{-\epsilon}\beta_{\tilde{\lambda}}\cdot\mathscr{R} + \nabla_\mu(\widetilde{\mathscr{Z}^\mu} + J^\mu_\Theta) - ((1+\gamma+S)\phi)\cdot\frac{\delta}{\delta\phi}S_0.$$

One need only thrice differentiate this equation to obtain the four point amplitude of $\nabla_\mu\mathcal{D}^\mu$. Note that on fixed points and cycles, where we will need this, the first term vanishes since $\widehat{B}_I = 0$. Also, the last term, which vanishes by the equations of motion, can be ignored for the computation of the amplitude. Most of the remaining terms vanish once the couplings are taken to be spacetime-independent (and the metric flat and $A_\mu = 0$). The remaining terms arise from the G and H^2

terms in $\beta_\lambda \cdot \mathscr{R}$. For a conformally flat metric, $\gamma_{\mu\nu} = \exp(-2\tau(x))\eta_{\mu\nu}$, one has (in d spacetime dimensions)

$$e^{-4\tau}G = 8(\partial^2\tau)^2 - 8\tau_{,\mu\nu}\tau^{,\mu\nu} - 16\tau_{,\mu}\tau_{,\nu}\tau^{,\mu\nu} - 8(d-3)\tau_{,\mu}\tau^{,\mu}\partial^2\tau + 2(d-1)(d-4)(\tau_{,\mu}\tau^{,\mu})^2$$
$$e^{-4\tau}H^2 = 4(\partial^2\tau)^2 - 4(d-2)\tau_{,\mu}\tau^{,\mu}\partial^2\tau + (d-2)^2(\tau_{,\mu}\tau^{,\mu})^2.$$

The cubic term in H^2 vanishes for an "on-shell" conformal factor $\partial^2\tau = 0$ and so the only contribution to the "on-shell" forward scattering amplitude is from G:

$$\mathcal{A}_{\rm fwd}(s)|_{\text{FP or cycle}} = -\beta_b(s^2+t^2+u^2)|_{t=0} = -2\beta_b s^2. \tag{5.1}$$

We assume there exists an RG trajectory from a UV fixed point or cycle to an IR fixed point or cycle. On this trajectory this equation no longer holds. However, we can inspect limiting behavior. Since $\mathcal{A}_{\rm fwd}/s^2$ depends on s only through the dimensionless ratio μ^2/s, its behavior is dictated by the renormalization group. Hence,

$$\lim_{s\to\infty}\frac{\mathcal{A}_{\rm fwd}(s)}{s^2} = \lim_{s\to\infty}\frac{\mathcal{A}_{\rm fwd}(s)|_{\text{FP or cycle}}}{s^2} = -2(\beta_b)_{\rm UV}$$

and

$$\lim_{s\to 0}\frac{\mathcal{A}_{\rm fwd}(s)}{s^2} = \lim_{s\to 0}\frac{\mathcal{A}_{\rm fwd}(s)|_{\text{FP or cycle}}}{s^2} = -2(\beta_b)_{\rm IR},$$

where $(\beta_b)_{\rm UV}$ and $(\beta_b)_{\rm IR}$ are the limiting UV and IR values of β_b along the trajectory and correspond to those on the fixed point or cycle. LPR do a careful analysis to validate the approach to these limiting values using conformal perturbation theory.

Following LPR we next consider the integral of $\mathcal{A}_{\rm fwd}(s)/s^3$ over the contour in Fig. 1. The integral over the semicircle I_1 cannot be easily computed, but in the limit that the radius of the semicircle vanishes it is reasonable that one can use the limiting value,



Fig. 1: The contour of integration for $\displaystyle\int \frac{ds}{s^3}\,\mathcal{A}_{\rm fwd}(s)$.

$$\int_{I_1}\frac{ds}{s^3}\mathcal{A}_{\rm fwd}(s) \approx \int_{I_1}\frac{ds}{s}2(\beta_b)_{\rm IR} = 2\pi i(\beta_b)_{\rm IR}, \tag{5.2}$$

where the last step corresponds to taking the vanishing limit of the radius of the semicircle I_1. Similarly, the large circle I_3 gives

$$\int_{I_3} \frac{ds}{s^3} \mathcal{A}_{\text{fwd}}(s) \approx \int_{I_3} \frac{ds}{s} 2(\beta_b)_{\text{UV}} = -2\pi i (\beta_b)_{\text{UV}}. \tag{5.3}$$

It follows from Cauchy's theorem that

$$\begin{aligned}(\beta_b)_{\text{UV}} - (\beta_b)_{\text{IR}} &= \frac{1}{2\pi i} \int_{I_2} \frac{ds}{s^3} \mathcal{A}_{\text{fwd}}(s) \\ &= \frac{1}{\pi} \int_0^\infty \frac{ds}{s^3} \operatorname{Im}(\mathcal{A}_{\text{fwd}}(s + i0)),\end{aligned}$$

where in the last line LPR assume crossing symmetry to write $\mathcal{A}_{\text{fwd}}(-s + i0) = \mathcal{A}^*_{\text{fwd}}(s + i0)$. Finally, the KS argument invokes the optical theorem that relates the imaginary part of the forward scattering amplitude to a positive-definite cross section to conclude that

$$(\beta_b)_{\text{UV}} - (\beta_b)_{\text{IR}} > 0.$$

We note in passing that the optical theorem is known to apply for forward scattering amplitudes of (on-shell) physical particles. It is not clear a priori that it applies to Green functions of composite operators at $p_i^2 = 0$, even if it corresponds to the scattering amplitude of would-be dilaton scattering. We think the assumption of positivity is reasonable, so we press on.

What steps in the argument above require special attention when the theory admits dimension-three currents? As we have pointed out, the dilatation current now has an additional J^μ term, but we have already accounted for this in the presentation above. Throughout the flow this makes no difference to the argument above, since the positivity of the integral over the segments I_3 of the contour follows from the optical theorem. For cycles one is not free to ignore the $\tau(x)$ dependence of the couplings or the background vector field in the anomaly equation. But on the cycle the couplings are covariantly constant. Hence, the terms that vanish at fixed points because of the constancy of couplings also vanish for cycles, but now because they are covariantly constant. Finally, the validity of the limits in (5.2) and (5.3) need to be established anew for limit cycles. However, the same method of conformal perturbation theory may now be applied, using instead scale-but-not-conformal perturbation theory to establish the result. Since it is only scaling that is used in this step of the argument by LPR, the proof goes through as presented there.

6. Conclusions

We have shown that a weak version of the c-theorem can be shown along the lines of the arguments presented by Cappelli, D'Appollonio, Guida and Magnoli and by Komargodski and Schwimmer for the more general case that a renormalization group flow goes from a fixed point or cycle to another fixed point or cycle. To this end we demonstrated that the quantity that plays the role of c is Jack and Osborn's $\widetilde{B}_b$ which is closely related to the a-anomaly (β_b in the notation of Jack and Osborn); these quantities agree at fixed points, but are not generally the same. We showed that given an RG trajectory linking fixed points and/or cycles, there is a non-RG trajectory linking them too on which $\widetilde{B}_b$ decreases monotonically from the UV to the IR.

We showed where the arguments presented by Luty, Polchinski and Rattazzi, that no renormalization-group limit cycles can exist in four-dimensional theories, are in error. In contrast with their results, we showed that β_b is constant on a cycle. To this end we reviewed (and presented a somewhat different derivation of) the consistency conditions of Jack and Osborn. The crucial observation is that Luty et al neglect to include the effect of essential counterterms.

We presented a calculation of the Lie-algebra function of coupling constants S introduced by Jack and Osborn. This is the first calculation of S to an order (third) in the loop expansion where it does not vanish. We then proved that $S = 0$ on fixed points and that S precisely corresponds to the generator Q of limit cycles when evaluated at any point on the limit cycle. This is a major improvement on the method of searching for limit cycles: one merely needs to find zeroes of the beta functions to the first order in the loop expansion (second order for Yang–Mills couplings) and evaluate S there. If $S = 0$ the zero corresponds to a fixed point, while if $S = Q \neq 0$ the zero corresponds to a limit cycle with Q the generator of the cycle.

Some questions remain which we intend to turn to in the future. Among them are:

- Are there renormalization group flows between fixed points and cycles?
- Are there models in four dimensions with limit cycles having tree-level bounded scalar potential?
- Are there limit cycles in supersymmetric four-dimensional field theories?

We look forward to answering these questions.

Acknowledgments

We would like to thank David Pirtskhalava for numerous discussions and for his comments on the manuscript. This material is based upon work supported by the US Department of Energy under contract DOE-FG03-97ER40546 and in part by the US National Science Foundation under Grant No. 1066293 and the hospitality of the Aspen Center for Physics.

Appendix A. The relation between the metrics χ^a_{ij} and χ^g_{ij}

The coefficient c_g of LPR appears to play the role of the "metric" χ^g_{ij} in the consistency condition (2.13). As we mention in the end of Section 2.2 and elaborate on further here, this is not the case. To see the connection with the work of JO, following LPR we write

$$\Delta W_{\text{anomaly}} = \frac{1}{2}\int d^4x\, d^4y\, \tau(x)\tau(y)\langle\Theta(x)\Theta(y)\rangle,$$

where $\Theta = \beta^i\mathcal{O}_i$, and therefore

$$\frac{d}{dt}\Delta W_{\text{anomaly}} = \frac{1}{2}\int d^4x\, d^4y\, \tau(x)\tau(y)\frac{d}{dt}\langle\Theta(x)\Theta(y)\rangle. \tag{A.1}$$

In Ref. [9, Eq. (3.18b)] Osborn finds the RGE for the product of two local renormalized operators,

$$-\frac{d}{dt}\langle\mathcal{O}_i(x)\mathcal{O}_j(0)\rangle + \partial_i\beta^j\langle\mathcal{O}_i(x)\mathcal{O}_j(0)\rangle + \partial_j\beta^k\langle\mathcal{O}_i(x)\mathcal{O}_j(0)\rangle = -\chi^a_{ij}\partial^2\partial^2\delta^{(4)}(x).$$

The quantity χ^a_{ij} can be thought of as the beta function associated with the counterterm needed in order to renormalize the correlator $\langle\mathcal{O}_i(x)\mathcal{O}_j(0)\rangle$. Because of this, it is negative-definite. Now since $-d\beta^i/dt = \beta^j\partial_j\beta^i$, it is easy to see that

$$\frac{d}{dt}\langle\Theta(x)\Theta(0)\rangle = \chi^a_{ij}\beta^i\beta^j\partial^2\partial^2\delta^{(4)}(x). \tag{A.2}$$

Using this in (A.1) we see that the metric of LPR is $-2\chi^a$, which is always positive. This suggests the question: is there a relation between χ^g and χ^a?

In the specific example of a gauge theory with a simple gauge group G and charged fermions in some representation, JO give [8, Eqs. (5.12)], at two loops,

$$\chi^a = -\frac{1}{2}\chi^g = -\frac{n_V}{8\pi^2 g^2}\left[1 + \left(17C - \frac{20}{3}R\right)h\right], \qquad h \equiv \frac{g^2}{16\pi^2}, \tag{A.3}$$

where $\mathrm{tr}(t^a_{\text{adj}}t^b_{\text{adj}}) = C\delta^{ab}$, R is similarly defined for the representation of the fermions, and $n_V = \dim(\text{Adj})$ is the number of vectors. However, the relation $\chi^g = -2\chi^a$ of (A.3) does not hold in general, and so the task of computing χ^g is complicated. Nevertheless, Weyl consistency conditions give the general relation between χ^a and χ^g [8, Eq. (3.23)]:

$$\chi^g_{ij} + 2\chi^a_{ij} - \bar{\chi}^a_{ijk}\beta^k = -\beta^k\partial_k V_{ij} - \partial_i\beta^j\, V_{kj} - \partial_j\beta^k\, V_{ik}, \tag{A.4}$$

where

$$\zeta V_{ij} = \bar{\chi}^a_{ijk}k^k g^k \qquad \text{(no sum over } k\text{)},$$

and

$$\bar{\chi}^a_{ijk} = \partial_k\chi^a_{ij} - \tfrac{1}{2}(\chi^b_{ikj} + \chi^b_{jki}),$$

with χ^b_{ijk} necessary to regulate infinities in three-point functions, and ζ defined as an operator counting the number of loops, whose form can be read off from $\mathcal{O}(\epsilon)$ terms of the finiteness condition (3.9e) of JO:

$$\zeta V_{ij} = (1 + k^k g^k\partial_k)V_{ij} + 2k^i V_{ij} \qquad \text{(no sum over the index carried by } k\text{)},$$

(cf. [8, Eq. (3.16b)]).

In our gauge-theory example we have

$$\chi^g + 2\chi^a - \bar{\chi}^a \beta^g = -\beta^g \frac{\partial V}{\partial g} - 2\frac{\partial \beta^g}{\partial g} V, \qquad \zeta V = \tfrac{1}{2}\bar{\chi}^a g, \tag{A.5}$$

where $\zeta V = (2 + \frac{1}{2} g\, \partial/\partial g)V = (2 + h\, \partial/\partial h)V$, the beta function for the gauge coupling is

$$\frac{1}{g}\beta^g = -\beta_0 h - \beta_1 h^2 + \mathcal{O}(h^3), \qquad \beta_0 = \frac{1}{3}(11C - 3R),$$

and

$$\bar{\chi}^a = \frac{\partial \chi^a}{\partial g} - \chi^b,$$

where χ^b is given at two loops by

$$\chi^{b(2)} = \frac{n_V}{4\pi^2 g^3}\left[1 + \frac{4}{3}(11C - 4R)h\right].$$

It follows that

$$\bar{\chi}^a = -\frac{n_V}{3\pi^2 g^3}[(11C - 4R)h + \mathcal{O}(h^2)].$$

Expanding $V = v_0 + v_1 h + \cdots$ gives $\zeta V = 2v_0 + 3v_1 h + \cdots = \frac{1}{2} g \bar{\chi}^a$, or

$$V = -\frac{n_V}{192\pi^4}(11C - 4R) + \mathcal{O}(h).$$

With this, (A.5) gives

$$\chi^g + 2\chi^a = -\frac{n_V \beta_0}{96\pi^4}\,(11C - 4R)\,h + \mathcal{O}(h^2), \tag{A.6}$$

and, therefore, beyond two-loop order, $\chi^g \neq -2\chi^a$.

To summarize, as shown in JO the negative-definiteness of χ^a_{ij} follows from the two point function of the operators $\mathcal{O}_i$ and holds generally. In general $\chi^g_{ij} + 2\chi^a_{ij}$ fails to vanish beyond the first few orders in the loop expansion. The positivity of χ^g_{ij} may fail non-perturbatively (for example, if its perturbative expression has finite radius of convergence).

Appendix B. How to calculate N_I and S

The calculation of JO's N_I proceeds order by order in perturbation theory. In this appendix we calculate contributions to N_I in a quantum field theory with real scalars and Weyl spinors up to two loops, and we also perform a three-loop calculation of the part of N_I that is needed in order to compute S.

As can be seen from (3.1), in order to calculate N_I we need to compute self-energies of scalars but with coupling constants as spectator fields. Equivalently, the calculation can be done by considering scalar self-energy diagrams and letting momentum come in from external legs and go out through couplings. From these diagrams we can then pick up the contribution linear in the momentum of the field and linear in the momentum of the coupling. After we antisymmetrize, we have a contribution to N_I.

It is perhaps helpful to remind the reader here that in a theory with scalars and fermions the I index can be either $(abcd)$ or $(a|ij)$. Let us also remark that S appears first at three loops in a theory with scalars and spinors. The reason is easily seen from (3.7): a diagram that contributes to N will only contribute to S if it is not symmetric under $a \leftrightarrow b$. As it turns out there are no such diagrams in scalar self-energies at one and two loops, but there are two such diagrams at three loops. Consequently, even if the theory contains gauge fields, diagrams with gauge fields will not contribute to S at three loops, but certainly will do so at higher order. Therefore, even in a gauge theory we don't need to include gauge fields in our leading-order calculation of S.

B.1. One loop

At one loop the calculation proceeds with no subtleties since renormalization is straightforward. The two diagrams that contribute to N_I and their corresponding counterterms are shown in Fig. 2.



Fig. 2: Diagrams that contribute to $N_{a|ij}$ at one loop and their corresponding counterterms.

A straightforward calculation gives

$$(N_{c|ij})_{ab} = -\frac{1}{16\pi^2\epsilon}\,\frac{1}{2}(y^*_{a|ij}\delta_{bc} - y^*_{b|ij}\delta_{ac}) + \text{finite},$$

and there is of course a complex conjugate $(N^*_{c|ij})_{ab}$.

In order to simplify the notation we write the result for the residue of the simple ϵ-pole in N_I in the form

$$16\pi^2(N^1_I)_{ab}\partial^\mu g_I = -\tfrac{1}{2}[\mathrm{tr}(y_a\partial^\mu y^*_b) + \text{h.c.} - \{a \leftrightarrow b\}],$$

where g_I on the left-hand side stands here for $y_{c|ij}$ or $y^*_{c|ij}$. Selecting the appropriate derivatives one easily reads off the corresponding N^1_I. Our result reproduces JO's equation (7.16) for ρ_I when we use Dirac spinors.

B.2. Two loops

At two loops there are three Feynman diagrams that contribute to N_I, listed in Fig. 3. The



Fig. 3: Feynman diagrams that contribute to N_I at two loops.

calculation of the residues of the simple ϵ-poles of N_I requires now a subtraction of subdivergences, something that proceeds, for the most part, in the usual way. However, there is a small subtlety, not seen in the usual treatments of renormalization, that we would like to point out. Clearly, the two right-most diagrams of Fig. 3 have subdivergences so we have to add to them the diagrams with the insertions of the corresponding counterterms. For the right-most diagram the graph with the insertion of the counterterm is



Now, when the momentum that comes in from, say the left external leg, flows out through the counterterm, then there are two diagrams that contribute, namely



and



where the momentum exits to the north-east or to the north-west depending on which vertex it flows out of in the original diagram in Fig. 3. In both cases the counterterm is the same, but the diagram with the insertion of the counterterm is different as a result of the difference in the momentum of the internal leg that the counterterm picks up. That is, had we retained different momenta for the various vertices, there would be two momenta associated with the counterterm.

The two loop result for N_I^1, previously unpublished, is

$$\begin{aligned}(16\pi^2)^2(N_I^1)_{ab}\partial^\mu g_I = &-\tfrac{1}{24}\lambda_{acde}\,\partial^\mu\lambda_{bcde} + [\tfrac{1}{4}\,\mathrm{tr}(y_a^* y_c \partial^\mu y_b^* y_c) + \tfrac{3}{8}\,\mathrm{tr}(y_a^*\partial^\mu y_c y_c^* y_b) \\ &+ \tfrac{1}{4}\,\mathrm{tr}(y_a^* y_c \partial^\mu y_c^* y_b) + \tfrac{3}{8}\,\mathrm{tr}(y_a^* y_c y_c^* \partial^\mu y_b) + \text{h.c.}] - \{a \leftrightarrow b\}.\end{aligned}$$

It follows that S vanishes at this order. This can be seen, term by term (when anti-symmetrized in a and b) by replacing g_I for $\partial^\mu g_I$.

B.3. Three loops

At three loops there are many diagrams that contribute to N_I, but only four are not symmetric under $a \leftrightarrow b$ and thus end up contributing to S. These diagrams are shown in Fig. 4, and we here only compute their contributions to N_I^1.



Fig. 4: Diagrams that contribute to N_I not symmetric under $a \leftrightarrow b$ and hence contribute to S at three loops.

From these diagrams (and the corresponding counterterms), using the methods for the calculation of three-loop integrals given in [13], we find

$$\begin{aligned}(16\pi^2)^3(N_I^1)_{ab}\partial^\mu g_I \supset & -\tfrac{1}{2}\operatorname{tr}(y_a\partial^\mu y_c^* y_d y_e^*)\lambda_{bcde} - \tfrac{1}{3}\operatorname{tr}(y_a y_c^* \partial^\mu y_d y_e^*)\lambda_{bcde} - \tfrac{1}{2}\operatorname{tr}(y_a y_c^* y_d \partial^\mu y_e^*)\lambda_{bcde}\\
& - \tfrac{5}{24}\operatorname{tr}(y_a y_c^* y_d y_e^*)\partial^\mu\lambda_{bcde} - \tfrac{1}{24}\operatorname{tr}(y_b\partial^\mu y_c^* y_d y_e^*)\lambda_{acde} - \tfrac{5}{24}\operatorname{tr}(y_b y_c^*\partial^\mu y_d y_e^*)\lambda_{acde}\\
& - \tfrac{1}{24}\operatorname{tr}(y_b y_c^* y_d\partial^\mu y_e^*)\lambda_{acde} - \tfrac{5}{24}\operatorname{tr}(\partial^\mu y_b y_c^* y_d y_e^*)\lambda_{acde} - \tfrac{7}{32}\operatorname{tr}(y_a\partial^\mu y_c^* y_d y_d^* y_b y_c^*)\\
& - \tfrac{7}{96}\operatorname{tr}(y_a y_c^*\partial^\mu y_d y_d^* y_b y_c^*) - \tfrac{23}{96}\operatorname{tr}(y_a y_c^* y_d\partial^\mu y_d^* y_b y_c^*) - \tfrac{7}{96}\operatorname{tr}(y_a y_c^* y_d y_d^*\partial^\mu y_b y_c^*)\\
& - \tfrac{7}{32}\operatorname{tr}(y_a y_c^* y_d y_d^* y_b\partial^\mu y_c^*) + \tfrac{1}{16}\operatorname{tr}(y_a\partial^\mu y_c^* y_c y_d^* y_b y_d^*) - \tfrac{5}{48}\operatorname{tr}(y_a y_c^*\partial^\mu y_c y_d^* y_b y_d^*)\\
& - \tfrac{1}{48}\operatorname{tr}(y_a y_c^* y_c\partial^\mu y_d^* y_b y_d^*) - \tfrac{7}{96}\operatorname{tr}(y_a y_c^* y_c y_d^*\partial^\mu y_b y_d^*) + \tfrac{1}{16}\operatorname{tr}(y_a y_c^* y_c y_d^* y_b\partial^\mu y_d^*)\\
& + \text{h.c.} - \{a \leftrightarrow b\},\end{aligned}$$

and since

$$S \equiv -k_I N_I^1 g_I = -N_{abcd}^1\lambda_{abcd} - (\tfrac{1}{2}N_{a|ij}^1 y_{a|ij} + \text{h.c.})$$

we finally obtain

$$(16\pi^2)^3 S_{ab} = \tfrac{5}{8}\operatorname{tr}(y_a y_c^* y_d y_e^*)\lambda_{bcde} + \tfrac{3}{8}\operatorname{tr}(y_a y_c^* y_d y_d^* y_b y_c^*) + \text{h.c.} - \{a \leftrightarrow b\}.$$

As already remarked in the main body, evaluating this on points in coupling space where we have found fixed points and cycles in Refs. [3, 5, 6], we find that S vanishes at all fixed points and equals Q on all cycles.

References

[1] Z. Komargodski & A. Schwimmer, *"On Renormalization Group Flows in Four Dimensions"*, JHEP **1112**, 099 (2011), `arXiv:1107.3987`.

[2] A. Cappelli, G. D'Appollonio, R. Guida & N. Magnoli, *"On the c theorem in more than two-dimensions"*, JHEP **1112**, (2000), `hep-th/0009119`. ❖ A. Cappelli, R. Guida & N. Magnoli, *"Exact consequences of the trace anomaly in four-dimensions"*, Nucl.Phys. **B618**, 371 (2001), `hep-th/0103237`.

[3] J.F. Fortin, B. Grinstein & A. Stergiou, *"Scale without Conformal Invariance: An Example"*, Phys.Lett. **B704**, 74 (2011), `arXiv:1106.2540`.

[4] J.F. Fortin, B. Grinstein & A. Stergiou, *"Scale without Conformal Invariance: Theoretical Foundations"*, JHEP **1207**, 025 (2012), `arXiv:1107.3840`. ❖ J.F. Fortin, B. Grinstein & A. Stergiou, *"Cyclic unparticle physics"*, Phys.Lett. **B709**, 408 (2012), `arXiv:1110.1634`.

[5] J.F. Fortin, B. Grinstein & A. Stergiou, *"Scale without Conformal Invariance at Three Loops"*, `arXiv:1202.4757`.

[6] J.F. Fortin, B. Grinstein & A. Stergiou, *"Scale without Conformal Invariance in Four Dimensions"*, `arXiv:1206.2921`.

[7] M.A. Luty, J. Polchinski & R. Rattazzi, *"The a-theorem and the Asymptotics of 4D Quantum Field Theory"*, `arXiv:1204.5221`.

[8] I. Jack & H. Osborn, *"Analogs for the c theorem for four-dimensional renormalizable field theories"*, Nucl.Phys. **B343**, 647 (1990).

[9] H. Osborn, *"Weyl consistency conditions and a local renormalization group equation for general renormalizable field theories"*, Nucl.Phys. **B363**, 486 (1991).

[10] J. Wess & B. Zumino, *"Consequences of anomalous Ward identities"*, Phys.Lett. **B37**, 95 (1971).

[11] Z. Komargodski, *"The Constraints of Conformal Symmetry on RG Flows"*, `arXiv:1112.4538`.

[12] A. Iorio, L. O'Raifeartaigh, I. Sachs & C. Wiesendanger, *"Weyl gauging and conformal invariance"*, Nucl.Phys. **B495**, 433 (1997), `hep-th/9607110`.

[13] K.G. Chetyrkin, M. Misiak & M. Münz, *"Beta functions and anomalous dimensions up to three loops"*, Nucl.Phys. **B518**, 473 (1998), `hep-ph/9711266`.